

A minimal classical sequent calculus free of structural rules

DOMINIC HUGHES

Stanford University*

April 22, 2009

Gentzen's classical sequent calculus **LK** has explicit structural rules for contraction and weakening. They can be absorbed (in a right-sided formulation) by replacing the axiom $P, \neg P$ by $\Gamma, P, \neg P$ for any context Γ , and replacing the original disjunction rule with Γ, A, B implies $\Gamma, A \vee B$.

This paper presents a classical sequent calculus which is also free of contraction and weakening, but more symmetrically: both contraction and weakening are absorbed into conjunction, leaving the axiom rule intact. It uses a blended conjunction rule, combining the standard context-sharing and context-splitting rules: Γ, Δ, A and Γ, Σ, B implies $\Gamma, \Delta, \Sigma, A \wedge B$. We refer to this system **M** as *minimal sequent calculus*.

We prove a *minimality theorem* for the propositional fragment **Mp**: any propositional sequent calculus S (within a standard class of right-sided calculi) is complete *if and only if* S contains **Mp** (that is, each rule of **Mp** is derivable in S). Thus one can view **M** as a minimal complete core of Gentzen's **LK**.

1 Introduction

The following Gentzen-Schütte-Tait [Gen39, Sch50, Tai68] system, denoted **GS1p** in [TS96], is a standard right-sided formulation of the propositional fragment of Gentzen's classical sequent calculus **LK**:

System GS1p		
$\frac{}{P, \neg P}$	$\frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \ \&$	$\frac{\Gamma, A_i}{\Gamma, A_1 \vee A_2} \ \oplus_i$
$\frac{\Gamma}{\Gamma, A} \ \text{W}$	$\frac{\Gamma, A, A}{\Gamma, A} \ \text{C}$	

Here P ranges over propositional variables, A, A_i, B range over formulas, Γ ranges over disjoint unions of formulas, and comma denotes disjoint union.¹ By defining a sequent as a disjoint union of formulas, rather than an ordered list, we avoid an exchange/permutation

*Visiting Scholar, Concurrency Group, Computer Science Department, Stanford University. I gratefully acknowledge my host, Vaughan Pratt.

¹We label the conjunction and disjunction rules with $\&$ and $\oplus$ for reasons which will become apparent later.

rule (cf. [TS96, §1.1]). Negation is primitive on propositional variables P , and extends to compound formulas by de Morgan duality.²

The structural rules, weakening W and contraction C , are absorbed in the following variant, a right-sided formulation of the propositional part of the calculus of [Ket44], called **GS3p** in [TS96].³

System GS3p		
$\frac{}{\Gamma, P, \neg P}$	$\frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \&$	$\frac{\Gamma, A, B}{\Gamma, A \vee B} \wp$

The new axiom $\Gamma, P, \neg P$ amounts to the original axiom $P, \neg P$ followed immediately by weakenings. This paper presents a propositional classical sequent calculus **Mp** which is also free of structural rules:

System Mp			
$\frac{}{P, \neg P}$	$\frac{\Gamma, \Delta, A \quad \Gamma, \Sigma, B}{\Gamma, \Delta, \Sigma, A \wedge B} \wedge$	$\frac{\Gamma, A, B}{\Gamma, A \vee B} \wp$	$\frac{\Gamma, A_i}{\Gamma, A_1 \vee A_2} \oplus_i$

A distinguishing feature of **Mp** is the *blended conjunction rule*⁴

$$\frac{\Gamma, \Delta, A \quad \Gamma, \Sigma, B}{\Gamma, \Delta, \Sigma, A \wedge B} \wedge$$

which combines the standard context-sharing and context-splitting conjunction rules:

$$\frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \& \quad \frac{\Delta, A \quad \Sigma, B}{\Delta, \Sigma, A \wedge B} \otimes$$

We refer to **Mp** as (cut-free propositional) *minimal sequent calculus*. In contrast to **GS3p**, contraction and weakening are absorbed symmetrically: both are absorbed into the conjunction rule, leaving the axiom rule intact.

Mp is evidently sound, since each of its rules can be derived (encoded) in **GS1p**. Theorem 1 (page 4) is completeness for formulas: a formula is valid iff it is derivable in **Mp**.⁵

² $\neg(A \vee B) = (\neg A) \wedge (\neg B)$ and $\neg(A \wedge B) = (\neg A) \vee (\neg B)$.

³ We label the disjunction rule as $\wp$ to distinguish it from the disjunction rule $\oplus$ of **GS1p**. The notation is derived from linear logic [Gir87].

⁴ By analogy with **GS3** and **GS3p** in [TS96], we reserve the symbol **M** for a full system with quantifiers, and use **Mp** to denote the propositional system. Following [TS96], we treat cut separately. To maximise emphasis on the blended conjunction rule, we omit quantifiers and cut in this paper.

⁵ Completeness here refers specifically to formulas, not to sequents. Section 6 discusses completeness for sequents.

1.1 Minimality

The blended conjunction rule $\wedge$ is critical for the liberation from structural rules: Proposition 2 (page 6) shows that relaxing it to the union of the two standard conjunction rules $\&$ and $\otimes$ breaks completeness.⁶ The main theorem of the paper (page 7) formalises the sense in which **Mp** is a minimal complete core of classical sequent calculus:

Theorem 2: Minimality

A standard sequent calculus S is complete iff $S \sqsupseteq \mathbf{Mp}$.

Here $S \sqsupseteq T$ (“ S contains T ”) iff every rule of T is derivable in S , and a *standard sequent calculus* is any propositional sequent calculus with the axiom $\overline{P}, \neg \overline{P}$ and any subset of the following *standard rules*:

$$\begin{array}{ccc} \frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \& \quad \quad \quad \frac{\Gamma, A, B}{\Gamma, A \vee B} \wp \quad \quad \quad \frac{\Gamma}{\Gamma, A} \mathbf{W} \\[1em] \frac{\Delta, A \quad \Sigma, B}{\Delta, \Sigma, A \wedge B} \otimes \quad \quad \quad \frac{\Gamma, A_i}{\Gamma, A_1 \vee A_2} \oplus_i \quad \quad \quad \frac{\Gamma, A, A}{\Gamma, A} \mathbf{C} \end{array}$$

2 Notation and terminology

Formulas are built from literals (propositional variables $P, Q, R \dots$ and their formal complements $\overline{P}, \overline{Q}, \overline{R}, \dots$) by the binary connectives **and** $\wedge$ and **or** $\vee$. Define **negation** or **not** $\neg$ as an operation on formulas (rather than as a connective): $\neg P = \overline{P}$ and $\neg \overline{P} = P$ for all propositional variables P , with $\neg(A \wedge B) = (\neg A) \vee (\neg B)$ and $\neg(A \vee B) = (\neg A) \wedge (\neg B)$.

We identify a formula with its parse tree, a tree labelled with literals at the leaves and connectives at the internal vertices. A **sequent** is a non-empty disjoint union of formulas.⁷ Comma denotes disjoint union. Throughout the document, $P, Q, \dots$ range over propositional variables, $A, B, \dots$ over formulas, and $\Gamma, \Delta, \dots$ over (possibly empty) disjoint unions of formulas.

A formula A is **valid** if it evaluates to 1 under all possible 0/1-assignments of its propositional variables (with the usual interpretation of $\wedge$ and $\vee$ on $\{0, 1\}$). A sequent $A_1, \dots, A_n$ is valid iff the formula $A_1 \vee (A_2 \vee (\dots \vee (A_{n-1} \vee A_n) \dots))$ is valid. A **subsequent** of a sequent Γ is any result of deleting zero or more formulas from Γ ; if at least one formula is deleted, the result is a **proper subsequent**.

⁶In other words, if we remove the $\wedge$ rule and add both the $\&$ and the $\otimes$ rules, the resulting system fails to be complete. The formula $((P \wedge Q) \vee (\overline{Q} \wedge P)) \vee \overline{P}$ becomes undervivable (see the proof of Proposition 2, page 6).

⁷Thus a sequent is a particular kind of labelled forest. This foundational treatment of formulas and sequents as labelled trees and forests sidesteps the common problem of “formulas” versus “formula occurrences”: disjoint unions of graphs are well understood in graph theory [Bol02].

3 Completeness

THEOREM 1 (COMPLETENESS) *Every valid formula is derivable in **Mp**.*

The proof is via the following auxiliary definitions and lemmas.

A sequent is **minimally valid**, or simply **minimal**, if it is valid while no proper subsequent is valid. For example, the sequents $P, \neg P$ and $P \wedge Q, \overline{Q} \wedge P, \overline{P}$ are minimal, while $P, \neg P, Q$ is not.

LEMMA 1 *Every valid sequent contains a minimal subsequent.*

Proof. Immediate from the definition of minimality. □

LEMMA 2 *Suppose a sequent Γ is a disjoint union of literals (i.e., Γ contains no $\wedge$ or $\vee$). Then Γ is minimal iff $\Gamma = P, \neg P$ for some propositional variable P .*

Proof. By definition of validity in terms of valuations, Γ is valid iff it contains a complementary pair of literals, i.e., iff $\Gamma = P, \neg P, \Delta$ with Δ a disjoint union of zero or more literals. Since $P, \neg P$ is valid, Γ is minimal iff Δ is empty. □

Suppose Γ and Δ are each disjoint unions of formulas (so each is either a sequent or empty). Write $\Gamma \subseteq \Delta$ if Γ results from deleting zero or more formulas from Δ .

LEMMA 3 *Suppose $\Gamma, A_1 \wedge A_2$ is minimal. Choose $\Gamma_1 \subseteq \Gamma$ and $\Gamma_2 \subseteq \Gamma$ such that Γ_1, A_1 and Γ_2, A_2 are minimal (existing by Lemma 1, since Γ, A_1 and Γ, A_2 are valid). Then every formula of Γ is in at least one of the Γ_i .*

Proof. Suppose the formula B of Γ is in neither Γ_i . Let Γ' be the result of deleting B from Γ . Then $\Gamma', A_1 \wedge A_2$ is a valid proper subsequent of $\Gamma, A_1 \wedge A_2$, contradicting minimality. (The sequent $\Gamma', A_1 \wedge A_2$ is valid since Γ_1, A_1 and Γ_2, A_2 are valid.) □

LEMMA 4 *Suppose $\Gamma, A \vee B$ is minimal and Γ, A is valid. Then Γ, A is minimal.*

Proof. If not, some proper subsequent Δ of Γ, A is valid. If Δ does not contain A , then it is also a proper subsequent of $\Gamma, A \vee B$, contradicting minimality. Otherwise let Δ' be the result of replacing A in Δ by $A \vee B$. Since Δ is valid, so also is Δ' . Thus Δ' is a valid proper subsequent of $\Gamma, A \vee B$, contradicting minimality. □

LEMMA 5 *Suppose $\Gamma, A \vee B$ is minimal and neither Γ, A nor Γ, B is valid. Then Γ, A, B is minimal.*

Proof. Suppose Γ, A, B had a valid proper subsequent Δ . Since neither Γ, A nor Γ, B is valid, Δ must contain both A and B . Let Δ' result from replacing A, B by $A \vee B$ in Δ . Then Δ' is a valid proper subsequent of $\Gamma, A \vee B$, contradicting minimality. □

Since a formula (viewed as a singleton sequent) is a minimal sequent, the Completeness Theorem (Theorem 1) is a special case of:

PROPOSITION 1 *Every minimal sequent is derivable in **Mp**.*

Proof. Suppose Γ is a minimal sequent. We proceed by induction on the number of connectives in Γ .

- *Induction base (no connective).* Since Γ is minimal, Lemma 2 implies $\Gamma = P, \neg P$, the conclusion of the axiom rule $\overline{P, \neg P}$.
- *Induction step (at least one connective).*

1. *Case:* $\Gamma = \Delta, A_1 \wedge A_2$. By Lemma 3, $\Gamma = \Sigma, \Delta_1, \Delta_2, A_1 \wedge A_2$ for Σ, Δ_1, A_1 and Σ, Δ_2, A_2 minimal. Write down the conjunction rule

$$\frac{\Sigma, \Delta_1, A_1 \quad \Sigma, \Delta_2, A_2}{\Sigma, \Delta_1, \Delta_2, A_1 \wedge A_2} \wedge$$

and appeal to induction with the two hypothesis sequents.

2. *Case:* $\Gamma = \Delta, A_1 \vee A_2$.

- (a) *Case:* Δ, A_i is valid for some $i \in \{1, 2\}$. Write down the disjunction rule

$$\frac{\Delta, A_i}{\Delta, A_1 \vee A_2} \oplus_i$$

then appeal to induction with Δ, A_i , which is minimal by Lemma 4.

- (b) *Case:* Δ, A_i is not valid for each $i \in \{1, 2\}$. Thus Δ, A_1, A_2 is minimal, by Lemma 5. Write down the disjunction rule

$$\frac{\Delta, A_1, A_2}{\Delta, A_1 \vee A_2} \wp$$

then appeal to induction with Δ, A_1, A_2 .

(Γ may match both 1 and 2 in the inductive step, permitting some choice in the construction of the derivation. There is choice in case 2(a) if both Δ, A_1 and Δ, A_2 are valid.) $\square$

Note that completeness does not hold for arbitrary valid sequents. For example, the sequent $P, \neg P, Q$ is valid but not derivable in **Mp**. A sequent is valid iff some some subsequence is derivable in **Mp**. Thus **Mp** is complete for sequents modulo final weakenings. In this sense, **Mp** is akin to system **GS5p** of [TS96, §7.4] (related to resolution). (See also Section 6.)

4 The Minimality Theorem

Relaxing blended conjunction to the pair of standard conjunction rules (context-sharing $\&$ and context-splitting $\otimes$) breaks completeness. Let **Mp**[−] be the following subsystem of **Mp**.⁸

⁸This precursor of **Mp** is (cut-free) multiplicative-additive linear logic [Gir87] with *tensor* $\otimes$ and *with* $\&$ collapsed to $\wedge$, and *plus* $\oplus$ and *par* $\wp$ collapsed to $\vee$.

System $\mathbf{Mp}^-$

$$\begin{array}{c}
 \frac{}{P, \neg P} \\
 \frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \& \quad \frac{\Gamma, A, B}{\Gamma, A \vee B} \wp \\
 \frac{\Delta, A \quad \Sigma, B}{\Delta, \Sigma, A \wedge B} \otimes \quad \frac{\Gamma, A_i}{\Gamma, A_1 \vee A_2} \oplus_i
 \end{array}$$

PROPOSITION 2 *System $\mathbf{Mp}^-$ is incomplete.*

Proof. We show that the valid formula $A = ((P \wedge Q) \vee (\overline{Q} \wedge P)) \vee \overline{P}$ is not derivable in $\mathbf{Mp}^-$. The placement of the two outermost $\vee$ connectives forces the last two rules of a potential derivation to be disjunction rules. Since $P \wedge Q, \overline{Q} \wedge P, \overline{P}$ is minimal (no proper subsequent is valid), the two disjunction rules must be $\wp$ rather than $\oplus$:

$$\frac{\frac{P \wedge Q, \overline{Q} \wedge P, \overline{P}}{(P \wedge Q) \vee (\overline{Q} \wedge P), \overline{P}} \wp}{((P \wedge Q) \vee (\overline{Q} \wedge P)) \vee \overline{P}} \wp$$

It remains to show that $P \wedge Q, \overline{Q} \wedge P, \overline{P}$ is not derivable in $\mathbf{Mp}^-$. There are only two connectives, both $\wedge$, so the last rule must be a conjunction.

1. *Case: the last rule is a context-sharing $\&$ -rule.*

(a) *Case: The last rule introduces $P \wedge Q$.*

$$\frac{P, \overline{Q} \wedge P, \overline{P} \quad Q, \overline{Q} \wedge P, \overline{P}}{P \wedge Q, \overline{Q} \wedge P, \overline{P}} \&$$

The left hypothesis $P, \overline{Q} \wedge P, \overline{P}$ cannot be derived in $\mathbf{Mp}^-$, since there is no Q to match the $\overline{Q}$ (and no weakening).

(b) *Case: The last rule introduces $\overline{Q} \wedge P$.* The same as the previous case, by symmetry, and exchanging $Q \leftrightarrow \overline{Q}$.

2. *Case: the last rule is a context-splitting $\otimes$ -rule.*

(a) *Case: The last rule introduces $P \wedge Q$.*

$$\frac{P, \Gamma \quad Q, \Delta}{P \wedge Q, \overline{Q} \wedge P, \overline{P}} \otimes$$

We must allocate each of $\overline{Q} \wedge P$ and $\overline{P}$ either to Γ or to Δ . If $\overline{Q} \wedge P$ is in Γ , then P, Γ is not derivable in $\mathbf{Mp}^-$, since it contains no Q to match the $\overline{Q}$. So $\overline{Q} \wedge P$ is in Δ . But then the $\overline{P}$ is required in both Γ and Δ .

- (b) *Case: The last rule introduces $\overline{Q} \wedge P$.* The same as the previous case, by symmetry, and exchanging $Q \leftrightarrow \overline{Q}$.

□

A **standard system** is any propositional sequent calculus containing the axiom $\overline{P}, \neg P$ and any of the following **standard rules**:

$$\begin{array}{ccc} \frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \& \quad \quad \quad \frac{\Gamma, A, B}{\Gamma, A \vee B} \wp \quad \quad \quad \frac{\Gamma}{\Gamma, A} W \\ \frac{\Delta, A \quad \Sigma, B}{\Delta, \Sigma, A \wedge B} \otimes \quad \quad \quad \frac{\Gamma, A_i}{\Gamma, A_1 \vee A_2} \oplus_i \quad \quad \quad \frac{\Gamma, A, A}{\Gamma, A} C \end{array}$$

Thus there are $2^6 = 64$ such systems (many of which will not be complete).

System S **contains** system T , denoted $S \sqsupseteq T$, if each rule of T is a derived rule of S . For example, system **GS1p** (page 1) contains **Mp** since the blended conjunction rule $\wedge$ and the disjunction rule $\wp$ of **Mp** can be derived in **GS1p**:

$$\begin{array}{ccc} \frac{\Gamma, \Delta, A \quad \Gamma, \Sigma, B}{\Gamma, \Delta, \Sigma, A \wedge B} \wedge & \longleftarrow & \frac{\frac{\Gamma, \Delta, A}{\Gamma, \Delta, \Sigma, A} W^* \quad \frac{\Gamma, \Sigma, B}{\Gamma, \Delta, \Sigma, B} W^*}{\Gamma, \Delta, \Sigma, A \wedge B} \& \\ \\ \frac{\Gamma, A, B}{\Gamma, A \vee B} \wp & \longleftarrow & \frac{\frac{\frac{\Gamma, A, B}{\Gamma, A, A \vee B} \oplus_2 \quad \frac{\Gamma, A, A \vee B}{\Gamma, A \vee B, A \vee B} \oplus_1}{\Gamma, A \vee B} C \end{array}$$

where W^* denotes a sequence of zero or more weakenings.

THEOREM 2 (MINIMALITY THEOREM) *A standard system is complete iff it contains **Mp**.*

4.1 Proof of the Minimality Theorem

Two systems are **equivalent** if each contains the other. For example, it is well known that **GS1p** (page 1) is equivalent to:⁹

$$\frac{}{P, \neg P} \quad \frac{\Delta, A \quad \Sigma, B}{\Delta, \Sigma, A \wedge B} \otimes \quad \frac{\Gamma, A_1, A_2}{\Gamma, A_1 \vee A_2} \wp \quad \frac{\Gamma}{\Gamma, A} W \quad \frac{\Gamma, A, A}{\Gamma, A} C$$

⁹This system is multiplicative linear logic [Gir87] plus contraction and weakening (with the connectives denoted $\wedge$ and $\vee$ instead of $\otimes$ and $\wp$).

via the following rule derivations:

$$\begin{array}{ccc}
\frac{\Delta, A \quad \Sigma, B}{\Delta, \Sigma, A \wedge B} \otimes & \longleftarrow & \frac{\frac{\Delta, A}{\Delta, \Sigma, A} W^* \quad \frac{\Sigma, B}{\Delta, \Sigma, B} W^*}{\Delta, \Sigma, A \wedge B} \& \\
\\
\frac{\Gamma, A_i}{\Gamma, A_1 \vee A_2} \oplus_i & \longleftarrow & \frac{\frac{\Gamma, A_i}{\Gamma, A_1, A_2} W}{\Gamma, A_1 \vee A_2} \wp \\
\\
\frac{\Gamma, A, B}{\Gamma, A \vee B} \wp & \longleftarrow & \frac{\frac{\frac{\Gamma, A, B}{\Gamma, A, A \vee B} \oplus_2 \quad \frac{\Gamma, A \vee B, A \vee B}{\Gamma, A \vee B} \oplus_1}{\Gamma, A \vee B} C \\
\\
\frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \& & \longleftarrow & \frac{\frac{\Gamma, A \quad \Gamma, B}{\Gamma, \Gamma, A \wedge B} \otimes}{\Gamma, A \wedge B} C^*
\end{array}$$

We shall abbreviate these four rule derivations as follows, and write analogous abbreviations for other rule derivations.¹⁰

$$\begin{array}{ccc}
\otimes & \longleftarrow & \& W \\
\oplus & \longleftarrow & \wp W \\
\& & \longleftarrow & \otimes C \\
\wp & \longleftarrow & \oplus C
\end{array}$$

4.1.1 The three complete standard systems

As a stepping stone towards the Minimality Theorem, we shall prove that, up to equivalence, there are only three complete standard systems.

We abbreviate a system by listing its non-axiom rules. For example, **GS1p** = (&, ⊕, W, C) and **Mp** = (∧, ⊕, ∖). Besides **GS1p**, we shall pay particular attention to the systems

$$\begin{array}{ll}
\mathbf{Pp} = (\otimes, \oplus, C) & \textit{Positive calculus} \\
\mathbf{Np} = (\&, \wp, W) & \textit{Negative calculus}
\end{array}$$

(Our terminology comes from polarity of connectives in linear logic [Gir87]: *tensor* ⊗ and *plus* ⊕ are positive, and *with* & and *par* ∖ are negative.)

¹⁰“Context-splitting conjunction ⊗ is derivable from context-sharing conjunction & and weakening W”, etc.

PROPOSITION 3 *Up to equivalence:*

- (1) **GS1p** = $(\&, \oplus, C, W)$ is the only complete standard system with both contraction C and weakening W ;
- (2) **Pp** = $(\otimes, \oplus, C)$ is the only complete standard system without weakening W ;
- (3) **Np** = $(\&, \wp, W)$ is the only complete standard system without contraction C .

The proof is via the following lemmas.

LEMMA 6 **Mp** = $(\wedge, \oplus, \wp)$ is contained in each of **Pp** = $(\otimes, \oplus, C)$, **Np** = $(\&, \wp, W)$ and **GS1p** = $(\&, \oplus, C, W)$.

Proof. **Pp** contains **Mp** since $\wedge \leftarrow C\otimes$,

$$\frac{\Gamma, \Delta, A \quad \Gamma, \Sigma, B}{\Gamma, \Delta, \Sigma, A \wedge B} \wedge \quad \leftarrow \quad \frac{\frac{\Gamma, \Delta, A \quad \Gamma, \Sigma, B}{\Gamma, \Gamma, \Delta, \Sigma, A \wedge B} \otimes}{\Gamma, \Delta, \Sigma, A \wedge B} C^*$$

(where C^* denotes zero or more consecutive contractions) and $\wp \leftarrow \oplus C$:

$$\frac{\Gamma, A, B}{\Gamma, A \vee B} \wp \quad \leftarrow \quad \frac{\frac{\frac{\Gamma, A, B}{\Gamma, A, A \vee B} \oplus_2}{\Gamma, A \vee B, A \vee B} \oplus_1}{\Gamma, A \vee B} C$$

Np contains **Mp** since $\wedge \leftarrow W\&$, and $\oplus \leftarrow W\wp$ (see page 8). **GS1p** = $(\&, \oplus, C, W)$ is equivalent to $(\otimes, \&, \oplus, \wp, C, W)$ since $\otimes$ and $\wp$ are derivable. Thus **GS1p** contains **Pp** (and **Np**), hence **Mp**. $\square$

LEMMA 7 **Pp** = $(C, \otimes, \oplus)$ and **Np** = $(\&, \wp, W)$ are complete.¹¹

Proof. Each contains **Mp** by Lemma 6, which is complete (Theorem 1). $\square$

LEMMA 8 *Up to equivalence, system **GS1p** = $(\&, \oplus, C, W)$ is the only complete standard system with both contraction C and weakening W .*

Proof. **GS1p** is complete (see e.g. [TS96], or by the fact that **GS1p** contains **Mp** which is complete). Any complete system must have a conjunction rule ($\otimes$ or $\&$) and a disjunction rule ($\oplus$ or $\wp$). In the presence of C and W , the two conjunctions are derivable from one other, as are the two disjunctions (see page 8). $\square$

LEMMA 9 *A complete standard system without weakening W must contain **Pp** = $(\otimes, \oplus, C)$.*

¹¹Recall that completeness refers to formulas, not sequents in general.

Proof. System $\mathbf{Mp}^- = (\otimes, \oplus, \&, \wp)$, with both conjunction rules and both disjunction rules, is incomplete (Proposition 2, page 6), therefore we must have contraction C.

Without the $\oplus$ rule, the valid formula $(P \vee \overline{P}) \vee Q$ is not derivable: the last rule must be $\wp$, leaving us to derive $P \vee \overline{P}, Q$, which is impossible without weakening W (i.e., with at most $\wp, \&, \otimes$ and C available), since, after a necessary axiom $\overline{P}, \overline{P}$ at the top of the derivation, there is no way to introduce the formula Q .

Without the context-splitting $\otimes$ rule, the valid formula $P \vee (Q \vee (\overline{P} \wedge \overline{Q}))$ is not derivable. The last two rules must be $\wp$, for if we use a $\oplus$ we will not be able to match complementary literals in the axioms at the top of the derivation. Thus we are left to derive $P, Q, \overline{P} \wedge \overline{Q}$, using $\&$ and C. The derivation must contain an axiom rule $\overline{P}, \overline{P}$. The next rule can only be a $\&$ (since $P, \overline{P}$ cannot be the hypothesis sequent of a contraction C rule). Since the only $\wedge$ -formula in the final concluding sequent $P, Q, \overline{P} \wedge \overline{Q}$ is $\overline{P} \wedge \overline{Q}$, and the $\&$ rule is context sharing, the $\&$ -rule must be

$$\frac{\overline{P}, \overline{P} \quad \begin{array}{c} \vdots \\ P, \overline{Q} \end{array}}{P, \overline{P} \wedge \overline{Q}} \&$$

but $P, \overline{Q}$ is not derivable. $\square$

LEMMA 10 *Up to equivalence, $\mathbf{Pp} = (\otimes, \oplus, C)$ is the only complete standard system without weakening W.*

Proof. By Lemma 7, $\mathbf{Pp}$ is complete. By Lemma 9, every W-free complete standard system contains $\mathbf{Pp}$. All other W-free standard systems containing $\mathbf{Pp}$ are equivalent to $\mathbf{Pp}$, since the standard rule derivations $\& \leftarrow \otimes C$ and $\wp \leftarrow \oplus C$ yield $\&$ and $\wp$ (see page 8). $\square$

LEMMA 11 *A complete standard system without contraction C must contain $\mathbf{Np} = (\&, \wp, W)$.*

Proof. System $\mathbf{Mp}^- = (\otimes, \oplus, \&, \wp)$, with both conjunction rules and both disjunction rules, is incomplete (Proposition 2, page 6), therefore we must have weakening W.

Without the $\wp$ rule, the valid formula $P \vee \overline{P}$ would not be derivable.

Without the $\&$ rule the valid formula $P \vee (\overline{P} \wedge \overline{P})$ would not be derivable. The last rule must be a $\wp$ (rather than a $\oplus$, otherwise we lack either P or $\overline{P}$), so we are left to derive $P, \overline{P} \wedge \overline{P}$. The last rule cannot be $\wp$ or $\oplus$, as the only connective is $\wedge$. It cannot be W, or else we lack either P or $\overline{P}$. It cannot be $\otimes$, as one of the two hypotheses will be the single formula $\overline{P}$. $\square$

LEMMA 12 *Up to equivalence, $\mathbf{Np} = (\&, \wp, W)$ is the only complete standard system without contraction C.*

Proof. By Lemma 7, $\mathbf{Np}$ is complete. By Lemma 11, every C-free complete standard system contains $\mathbf{Np}$. All other C-free standard systems containing $\mathbf{Np}$ are equivalent to $\mathbf{Np}$, since the standard rule derivations $\otimes \leftarrow \& W$ and $\oplus \leftarrow \wp W$ yield $\otimes$ and $\oplus$ (see page 8). $\square$

Proof of Proposition 3. Parts (1), (2) and (3) are Lemmas 8, 10 and 12, respectively. $\square$

LEMMA 13 *Every standard complete system has contraction C or weakening W .*

Proof. Otherwise it is contained in $\mathbf{Mp}^- = (\otimes, \oplus, \&, \wp)$, which is incomplete (Prop. 2). $\square$


THEOREM 3 *Up to equivalence, there are only three complete standard systems:*

1. *The Gentzen-Schütte-Tait system $\mathbf{GS1p} = (\&, \oplus, C, W)$.*
2. *Positive calculus $\mathbf{Pp} = (\otimes, \oplus, C)$.*
3. *Negative calculus $\mathbf{Np} = (\&, \wp, W)$.*

Proof. Proposition 3 and Lemma 13. $\square$

Proof of Minimality Theorem (Theorem 2). Each of the three complete standard systems contains $\mathbf{Mp}$ (Lemma 6). $\square$

The three inequivalent complete standard systems $\mathbf{GS1p}$, $\mathbf{Pp}$ and $\mathbf{Np}$, together with propositional minimal sequent calculus $\mathbf{Mp}$, sit in the following Hasse diagram of containments:



Thus we can view propositional minimal sequent calculus $\mathbf{Mp}$ as a minimal complete core of $\mathbf{GS1p}$, hence of (propositional) Gentzen's $\mathbf{LK}$.

5 Extended Minimality Theorem

Define an *extended system* as one containing the axiom rule $\overline{P, \overline{P}}$ and any of the following rules. (We have extended the definition of *standard system* by making blended conjunction available.)

Extended system rules

$$\begin{array}{c}
\frac{\Delta, A \quad \Sigma, B}{\Delta, \Sigma, A \wedge B} \otimes \quad \frac{\Gamma, \Delta, A \quad \Gamma, \Sigma, B}{\Gamma, \Delta, \Sigma, A \wedge B} \wedge \quad \frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \wedge B} \& \\
\\
\frac{\Gamma, A_i}{\Gamma, A_1 \vee A_2} \oplus_i \quad \frac{\Gamma, A, B}{\Gamma, A \vee B} \wp \\
\\
\frac{\Gamma, A, A}{\Gamma, A} C \quad \frac{\Gamma}{\Gamma, A} W
\end{array}$$

The Minimality Theorem (Theorem 2, page 7) extends as follows.

THEOREM 4 (EXTENDED MINIMALITY THEOREM) *An extended system is complete iff it contains propositional minimal sequent calculus **Mp**.*

To prove this theorem, we require two lemmas.

LEMMA 14 *Suppose S is a complete extended system with the blended conjunction rule $\wedge$, and with at least one of contraction C or weakening W . Then S is equivalent to a standard system.*

Proof. If S has weakening W , let S' be the result of replacing the blended conjunction rule $\wedge$ in S by context-sharing conjunction $\&$; otherwise S has contraction, and let S' result from replacing $\wedge$ by context-splitting $\otimes$. Then S' is equivalent to S , since $\wedge \leftarrow \otimes C$ (page 9) and $\wedge \leftarrow \& W$ (page 7). $\square$

LEMMA 15 *Suppose S is a complete extended system with neither contraction C nor weakening W . Then S is equivalent to propositional minimal sequent calculus **Mp**.*

Proof. Since $\mathbf{Mp}^- = (\otimes, \&, \oplus, \wp)$ is incomplete (Proposition 2, page 6), S must have the blended conjunction rule $\wedge$ either directly or as a derived rule. Since S is complete, it must have a disjunction rule, therefore it could only fail to be equivalent to $\mathbf{Mp} = (\wedge, \oplus, \wp)$ if (a) it has $\oplus$ and $\wp$ is not derivable, i.e., S is equivalent to $(\wedge, \oplus)$, or (b) it has $\wp$ and $\oplus$ is not derivable, i.e., S is equivalent to $(\wedge, \wp)$. In case (a), the valid formula $P \vee \overline{P}$ would not be derivable, and in case (b) the valid formula $(P \vee \overline{P}) \vee Q$ would not be derivable, either way contradicting the completeness of S . $\square$

Proof of the Extended Minimality Theorem (Theorem 4). Suppose S is a complete extended system. If S has contraction C or weakening W then it is equivalent to a standard system by

Lemma 14, hence contains **Mp** by the original Minimality Theorem. Otherwise S is equivalent to **Mp** by Lemma 15, hence in particular contains **Mp**.

Conversely, suppose S is an extended system containing **Mp**. Then S is complete since **Mp** is complete. $\square$

We also have the following extension of Theorem 3 (page 11), which stated that, up to equivalence, there are only three complete standard systems, **GS1p**, **Pp** and **Np**.

THEOREM 5 *Up to equivalence, there are only four complete extended systems:*

1. *The Gentzen-Schütte-Tait system **GS1p** = $(\&, \oplus, C, W)$.*
2. *Positive calculus **Pp** = $(\otimes, \oplus, C)$.*
3. *Negative calculus **Np** = $(\&, \wp, W)$.*
4. *Propositional minimal sequent calculus **Mp** = $(\wedge, \oplus, \wp)$.*

Proof. Theorem 3 together with Lemmas 14 and 15. $\square$

6 Degrees of completeness

We defined a system as *complete* if every valid formula (singleton sequent) is derivable. To avoid ambiguity with forthcoming definitions, let us refer to this default notion of completeness as **formula-completeness**. Define a system as **minimal-complete** if every minimal¹² sequent is derivable, and **sequent-complete** if every valid sequent is derivable. (Thus *sequent-complete* implies *minimal-complete* implies *formula-complete*.)

For a minimal-complete system S , a sequent Γ is valid iff a subsequence of Γ is derivable in S . Thus a minimal-complete system S can be viewed as sequent-complete, modulo final weakenings. (Cf. system **GS5p** of [TS96, §7.4] (related to resolution).)

PROPOSITION 4 **Pp** = $(\otimes, \oplus, C)$ and **Mp** = $(\wedge, \oplus, \wp)$ are formula-complete and minimal-complete, but not sequent-complete.

Proof. We have already proved that **Mp** (hence also **Pp**, by containment) is minimal-complete (Proposition 1).


We show that the valid (non-minimal) sequent $P, \overline{P}, Q$ is not derivable in **Pp** (hence also in **Mp**). A derivation must contain an axiom rule $\overline{P}, \overline{P}$. This cannot be followed by a $\otimes$ or $\oplus$ rule, otherwise we introduce a connective $\wedge$ or $\vee$ which cannot subsequently be removed by any other rule before the concluding sequent $P, \overline{P}, Q$. Neither can it be followed by contraction C , since there is nothing to contract. $\square$

PROPOSITION 5 **Np** = $(\&, \wp, W)$ is formula-, minimal- and sequent-complete.

¹²Recall that a valid sequent is minimal if no proper subsequence is valid.

Proof. **Np** is minimal-complete since it contains **Mp**. Suppose Γ is a valid but not minimal sequent. Choose a minimal subsequent Δ of Γ (see Lemma 1, page 4). By minimal-completeness, Δ has a derivation. Follow this with weakenings to obtain Γ . $\square$

Below we have annotated our Hasse diagram with completeness strengths.



7 Possible future work

1. *Cut*. Chapter 4 of [TS96] gives a detailed analysis of cut for Gentzen systems. One could pursue an analogous analysis of cut for minimal sequent calculus. Aside from context-splitting and context-sharing cut rules

$$\frac{\Delta, A \quad \Sigma, \neg A}{\Delta, \Sigma} \text{cut}_{\otimes} \qquad \frac{\Gamma, A \quad \Gamma, \neg A}{\Gamma} \text{cut}_{\&}$$

one might also investigate a blended cut rule:

$$\frac{\Gamma, \Delta, A \quad \Gamma, \Sigma, \neg A}{\Gamma, \Delta, \Sigma} \text{cut}$$

2. *Quantifiers*. Explore the various ways of adding quantifiers to **Mp**, for a full first-order system **M**.
3. *Mix (nullary multicut)*. Gentzen's multicut rule

$$\frac{\Delta, A_1, \dots, A_m \quad \Sigma, \neg A_1, \dots, \neg A_n}{\Delta, \Sigma}$$

in the nullary case $m = n = 0$ has been of particular interest to linear logicians [Gir87], who call it the *mix* rule. One could investigate context-splitting, context-sharing and blended incarnations:

$$\frac{\Delta \quad \Sigma}{\Delta, \Sigma} \text{mix}_{\otimes} \quad \frac{\Gamma \quad \Gamma}{\Gamma} \text{mix}_{\&} \quad \frac{\Gamma, \Delta \quad \Gamma, \Sigma}{\Gamma, \Delta, \Sigma} \text{mix}$$

References

- [Bol02] B. Bollobás. *Modern Graph Theory*. Springer-Verlag, 2002.
- [Gen39] G. Gentzen. Untersuchungen über das logische Schließen I, II. *Mathematische Zeitschrift*, 39:176–210 405–431, 1939. Translation in [Gen69, 68–131].
- [Gen69] G. Gentzen. *The Collected Papers of Gerhard Gentzen*. North-Holland, 1969. Edited and introduced by M.E. Szabo.
- [Gir87] J.-Y. Girard. Linear logic. *Theoretical Computer Science*, 50:1–102, 1987.
- [Ket44] J. Ketonen. Untersuchungen zum Prädikatenkalkül. *Annales Academiae Scientiarum Fennicae, ser. A, I. Mathematica-physica*, 23, 1944.
- [Sch50] K. Schütte. Schlußweisen-Kalküle der Prädikatenlogik. *Mathematische Annalen*, 123:166–186 405–431, 1950.
- [Tai68] W. W. Tait. Normal derivability in classical logic. In *The Syntax and Semantics of Infinitary Languages*, volume 72 of *Lecture Notes in Computer Science*, pages 204–236. Springer-Verlag, 1968.
- [TS96] A. S. Troelstra and H. Schwichtenberg. *Basic Proof Theory*. Cambridge Tracts in Theoretical Computer Science, Cambridge University Press, Cambridge, U.K., 1996.
- [Urb00] C. Urban. *Classical Logic and Computation*. Ph.D. thesis, University of Cambridge, 2000.
- [UB01] C. Urban and G. Bierman. Strong Normalisation of Cut-Elimination in Classical Logic. *Fundamenta Informaticae*, 45:123–155, 2001.